

COVID-19 VACCINE DOSAGE SCHEDULE

The below table summarizes the schedule of when additional doses and booster doses can be administered for individuals who **do not** have a weakened immune system.

Individuals **with** a weakened immune system should visit www.sedgwickcounty.org/covid-19/vaccine/additional-dosebooster-information/.

| Manufacturer* | Age Group | Total Number of Doses Recommended | Number of Primary and/or Additional Doses | Number of Booster Doses | Interval Between 1st and 2nd Dose | Interval Between 2nd and 3rd Dose | Interval Between 3rd and 4th Dose |
|-------------------|------------------|-----------------------------------|---|-------------------------|-----------------------------------|-----------------------------------|-----------------------------------|
| Pfizer | 6 months—4 years | 3 | 3 | N/A | 3—8 weeks | ≥ 8 weeks | N/A |
| Pfizer | 5—11 years | 3 | 2 | 1 | 3—8 weeks | ≥ 5 months | N/A |
| Pfizer | 12—49 years | 3 | 2 | 1 | 3—8 weeks | ≥ 5 months | N/A |
| Pfizer | 50 years + | 4 | 2 | 2 | 3—8 weeks | ≥ 5 months | ≥ 4 months |
| Moderna | 6 months—5 years | 2 | 2 | N/A | 4—8 weeks | N/A | N/A |
| Moderna | 6—17 years | 2 | 2 | N/A | 4—8 weeks | N/A | N/A |
| Moderna | 18—49 years | 3 | 2 | 1 | 4—8 weeks | ≥ 5 months | N/A |
| Moderna | 50 years + | 4 | 2 | 2 | 4—8 weeks | ≥ 5 months | ≥ 4 months |
| Johnson & Johnson | 18—49 years | 2 | 1 | 1 | ≥ 2 months | N/A | N/A |
| Johnson & Johnson | 50 years + | 3 | 1 | 2 | ≥ 2 months | ≥ 4 months | N/A |

* mRNA COVID-19 vaccines are preferred over Johnson & Johnson COVID-19 Vaccine for both primary and booster doses; an mRNA COVID-19 must be used for the second booster dose



Last updated 6/27/2022

PLEASE HELP PREVENT THE SPREAD OF COVID-19

