COVID-19 VACCINE DOSAGE SCHEDULE

The below table summarizes the schedule of when additional doses and booster doses can be administered for individuals who **do not** have a weakened immune system.

Individuals with a weakened immune system should visit www.sedgwickcounty.org/covid-19/vaccine/additional-dosebooster-information/.

Manufacturer*	Age Group	Total Number of Doses Recommended	Number of Primary and/or Additional Doses	Number of Booster Doses	Interval Between 1st and 2nd Dose	Interval Between 2nd and 3rd Dose
Pfizer	6 months— 4 years	3	3	N/A	3—8 weeks	≥ 8 weeks
Pfizer	5—11 years	3	2	1	3—8 weeks	≥ 2 months
Pfizer	12—17 years	3	2	1	3—8 weeks	≥ 2 months
Pfizer	18 years +	3	2	1	3—8 weeks	≥ 2 months
Moderna	6 months— 5 years	2	2	1	4—8 weeks	≥ 2 months
Moderna	6—17 years	3	2	1	4—8 weeks	≥ 2 months
Moderna	18 years +	3	2	1	4—8 weeks	≥ 2 months
Johnson & Johnson	18—49 years	2	1	1	≥ 2 months	N/A
Johnson & Johnson	50 years +	3	1	2	≥ 2 months	≥ 4 months
Novavax	12 years +	2	2	N/A	3—8 weeks	≥ 6 months**

^{*} mRNA COVID-19 vaccines are preferred over Johnson & Johnson COVID-19 Vaccine for both primary and booster doses; an mRNA COVID-19 must be used for the second booster dose

^{**} Individuals should receive the Novavax booster if they have completed a primary series at least 6 months prior, have not received previous boosters, and cannot or will not get the bivalent booster.



